

Review Article

Moringa Oleifera- A Wonder Plant for Male Fertility Preservation

Guruprasad N*, Sathish Kumar A**, Guruprasad K***

*Senior Research Fellow, **Professor and HOD, ***Professor, Department of Clinical Embryology, Kasturba Medical College, Manipal Academy of Higher Education, Manipal, Karnataka.



Guruprasad Nayak is working as Senior Research Fellow (SRF) and pursuing PhD at Department of Clinical Embryology, Kasturba Medical College (KMC), Manipal Academy of Higher Education (MAHE). He obtained his master (M.Sc.) degree in Medical Biochemistry from KMC, MAHE. He worked as research assistant in this institute for 3 years in ICMR funded project. He has published his research findings in reputed journals such as Andrologia, Journal of Ethnopharmacology, Environmental Toxicology and Pharmacology, Biomedicine and Pharmacotherapy, Journal of Assisted Reproduction and Genetics. His areas of interest are reproductive toxicology, proteomics and metabolomics.

Corresponding author - Dr.Guruprasad Kalthur (guru.kalthur@manipal.edu)

Chettinad Health City Medical Journal 2018; 7(3): 106 - 113

Abstract

Since time immemorial, *Moringa oleifera* Lam. (MO) is a highly valued plant with potential application in the area of nutrition and medicine. All parts of the plant have been used in preparing Ayurvedic medicines while the leaves and pods have been used in preparing traditional dishes. The leaves of this plant are most commonly used as they are rich in amino acids, vitamins, poly-phenols, flavonoids, alkaloids and phytosterols, which attribute to the nutritive and medicinal properties. The tissue and cyto-protective action of MO against various toxic agents is gaining tremendous importance in the field of toxicology. However, very few reports have highlighted the protective effect of MO and its active components against gonadotoxic agents. The present review mainly focuses on the medicinal properties of MO with special reference to its role in male fertility preservation.

Key words: Moringaoleifera, Chemotherapy, Testicular damage, Chemoprotection, Antioxidant

Introduction

Moringa oleifera (MO) is a fast growing tree belonging to monogeneric family Moringaceae, a native of India. The genus *Moringa* has 13 other species native to the African continent¹ and are distributed in tropical and subtropical regions of the world (Fig 1). The plant has been used from ancient times to prepare various dishes and herbal medicines owing to its rich nutrient contents and therapeutic values. Hence, it is popularly known as 'Miracle tree' or 'Nature gift' or 'Mothers best friend'. In India, MO is commonly referred to as 'drumstick tree', due to the shape of its fruit which resembles a drum stick. It is also called as 'horse radish tree' due to the pungent smell of its roots. The plant has a diverse application such as source of food, animal fodder, natural coagulants, forestry products, fertilizer, alley cropping and fueling.²



Fig 1: Global distribution of *Moringa* species (shown in green color)

Taxonomy of *Moringa oleifera* Lam.

Kingdom: Plantae
 Phylum: Tracheophyta
 Class: Magnoliopsida
 Order: Brassicales
 Family: Moringaceae
 Genus: *Moringa*
 Species: *oleifera*

Moringa oleifera plant as a nutritive source

Moringa oleifera is a highly nutritive plant. MO seeds are rich in monounsaturated fatty acids and proteins with sulfur containing amino acids.³ Interestingly, *Moringa* species grow in zones where malnutrition is more evident such as drought affected areas. The leaves, pods and seeds (Fig 2) of this plant are nutritious and most commonly used for preparing traditional dishes. In India, *M. oleifera* is commercially cultivated mainly for its pods as a vegetable⁴; the pods are rich in proteins, carbohydrates and dietary fibers.⁵ The leaves are rich in proteins, essential amino acids, vitamins, minerals and antioxidants which serve as natural remedy in preventing malnutrition in children of under-developed countries.⁶⁻⁸ A randomized blind placebo control study showed that MO leaf powder can improve nutritional intake and nutritional status in HIV infected patients undergoing anti-retroviral therapy thereby ensuring good immuno-metabolic response.⁹



Fig 2: Different parts of *Moringa oleifera* plant

Use of *Moringa oleifera* in Ayurvedic medicine

Moringa oleifera is the plant of choice for the preparation of various traditional medicines since ancient times. All parts of the plant have shown immense health benefits, particularly in improving function of reproductive, circulatory and immune system. Various studies have shown that MO and its active ingredients, either alone or as a multi-herbal preparation play a promising role in combating a wide range of pathologies including cancer, diabetes, ulcer, oxidative stress, infection and infertility (Table 1). Among different parts of the plant, leaves have been widely studied for its various therapeutic properties as detailed below.

a. Anticancer effect: Aqueous extract of MO leaves hinder the proliferation of alveolar cancer cells by inducing oxidative stress, DNA fragmentation and apoptosis. Moreover, the anti-proliferative effect of the extract was greater in cancer cells than normal cells¹⁰; thus it can be speculated that an

Parts	Composition	Medicinal property
Roots	Benzyl isothiocyanate Phenethyl isothiocyanate	Anticancer Neuro-protective Anti-fertility
Stem (Bark)		Antibacterial action Anticancer effect Insulin sensitization
Leaves	Glucomoringin isothiocyanate, Quercetin, Benzyl isothiocyanate, Vicenin-2, Chlorogenic acid, Gallic acid, Kaempferol, Rosmarinic acid and Rutin	Anticancer Chemo-sensitization
		Anti-diabetic
		Anti-bacterial Anti-viral
		Antianalgesic, anti-inflammatory and antioxidant Wound healing
Pods and seeds	Isothiocyanates	Anti-hypertension Antioxidant and antibacterial action Anti-inflammatory action Antiulcer
Flowers	Trypsin inhibitor	Larvicidal action on <i>Aedes aegypti</i> (dengue fever)

Table 1: Medicinal properties of different parts of *Moringa oleifera* plant

active principle of the extract specifically targets the cancer cells, thereby it can potentially decrease the side effects on normal tissues unlike many of the chemotherapeutic drugs. Aqueous extract of MO leaves increased the cytotoxic effect of chemotherapeutic agents on pancreatic cancer cells¹¹ by down-regulating nuclear factor-kappa beta (NF- κ B) indicating chemo-sensitization of the extract. Ethanolic extract of leaves exhibited anticancer activity in breast and colorectal cancer cell lines.¹² The crude extract and certain isolates of the plant such as glucomoringin isothiocyanate, quercetin and benzyl isothiocyanate have been shown to increase the apoptosis in cancer cell lines.¹³⁻¹⁶

- b. Antidiabetic effects: Methanolic extract of MO leaves prevented diabetes induced nephrotoxicity through its hyperglycemic, antioxidant and anti-inflammatory action.¹⁷ Aqueous extract of leaf showed potent anti-diabetic effect against streptozotocin-induced diabetes in rats.¹⁸ The study on type 2 diabetes-induced mice and rats revealed that the extract of MO exhibited hypoglycemic effect, ameliorated oxidative stress, renal and hepatic dysfunction and improved glycogen synthesis and modulated lipid metabolism.¹⁹⁻²² It has been demonstrated that MO leaf extract can mitigate alloxan-induced diabetes by regenerating β -cells and modulate the expression of pyruvate carboxylase.²³ Furthermore, the extract from bark showed improvement in insulin resistance induced by dexamethasone.²⁴
- c. Antimicrobial action: The water-soluble lectin extracted from MO exhibited antibacterial action against *Serratia marcescens* and *Bacillus* species by inducing cell wall damage.²⁵ The bark showed significant beneficial effect in the management of urinary tract infections.²⁶ The aqueous extract from MO leaves showed protection against HBV infection by reducing fibrosis markers, IL-6 and HBsAg secretion.²⁷ The flower extract inhibited the larval growth of *Aedes aegypti*, a vector of dengue fever, by targeting trypsin and acetyl choline esterase activity.²⁸
- d. Other properties: Aqueous extract of MO leaves contain phytochemicals like vicenin-2, chlorogenic acid, gallic acid, quercetin, kaempferol, rosmarinic acid and rutin which exhibited potential wound healing property.^{29,30} The methanolic extract showed analgesic, anti-inflammatory and antioxidant properties in animal models.³¹⁻³³ The active compounds in pods and seeds exhibited hypotensive and anti-ulcer properties respectively.^{34,35} The hydro-ethanolic extract of MO leaves showed cerebro-protective effect against ischemic stroke by decreasing oxidative stress³⁶ and penicillin induced convulsion, locomotor behavior by modulating the secretion of neurotransmitters.³⁷

Role of *Moringa oleifera* in tissue protection

The tissue protective action of MO is summarized in Table 2. The studies have shown that MO and its active components impart protective action against certain toxic chemicals and radiation. Aqueous extract of MO leaves showed amelioration of radiation-induced oxidative stress in mouse hepatocytes by inhibiting translocation of Nf-kb, decreasing lipid peroxidation and increasing the activity of antioxidant enzymes.³⁸ The toxic effect of radiation on bone marrow cell was mitigated by the extract by decreasing induction of chromosomal aberrations and micronucleus formation.³⁹ A recent study revealed that butyl p-hydroxyphenyl-acetate (MIMO₂), a novel compound isolated from MO leaves prevented oxidative stress and DNA damage induced by a metal vanadium, a potent neuro-toxicant.⁴⁰ Further, MO leaf extract showed nephro-protective effect against acetaminophen, a common analgesic and antipyretic by modulating activity of antioxidant enzymes and anti-inflammatory molecules.⁴¹ Similarly, another study exhibited the mitigating effect of MO leaves on acetaminophen induced hepatotoxicity by restoring glutathione level.⁴² The ethanolic extract of MO leaves showed ameliorating effect against anti-tuberculosis drugs (isoniazid, rifampicin, and pyrazinamide) - induced hepatotoxicity.⁴³ The oil extracted from MO

seeds showed protective effect against carbon tetrachloride (CCl₄)-induced hepatitis by decreasing lipid peroxidation and antioxidant status in rat hepatocytes.⁴⁴

Protective role of *Moringa oleifera* in male gonadal toxicity

It is known that certain chemicals and radiation are highly toxic to testes as they affect structure and/or functioning of testes. In literature, there are very few reports on protective action of MO against testicular toxicity. The ethanolic extract of leaves showed the mitigating effect against chromium-induced testicular toxicity and improvement in sperm functions by preventing oxidative stress and increasing testosterone in rats.⁴⁵ The oil produced from MO, mitigated mercury induced testicular toxicity by improving steroidogenesis and antioxidant status in rats.⁴⁶ However, there are no studies on the protective role of MO against chemotherapy and radiotherapy; which are known to cause severe damaging effects on testes imparting infertility. Therefore, we have conducted experiments to assess the testicular protective effect of ethanolic extract of MO leaves against cyclophosphamide, a broad spectrum anticancer drug.

Role of *Moringa oleifera* in Fertility Preservation

In recent years, the number of cancer survivors have been increasing due to the tremendous advancement in diagnostic and therapeutic strategies in cancer treatment. However, due to the non-specific action of radiation and anticancer drugs, these agents can cause toxicity in normal cells leading to various long-term health issues. Testis is highly vulnerable to these toxic agents due to the presence of rapidly proliferating spermatogonial stem cells (SSC). Depletion of SSC leads to temporary or permanent arrest in spermatogenesis. Therefore, cancer survivors have compromised fertility potential which in turn affects the quality of life post chemotherapy and radiotherapy.

Fertility preservation is a promising option to preserve reproductive capacity of cancer patients undergoing chemo- and radio- therapies. The currently available methods of fertility preservation are depicted in Fig 3.

MO Extract / Phytochemicals	Toxic agents	Tissue/ cell
Leaf extract	Radiation	Liver
		Bone marrow
Leaf extract	Acetaminophen	Kidney
		Liver
Leaf extract	Antituberculosis drugs- isoniazid, rifampicin, and pyrazinamide	Liver
Leaf extract	Chromium	Testis
Leaf extract	Cyclophosphamide	
Butyl p-hydroxy phenyl-acetate	Vanadium	Neurons

Table 2: Tissue protective action of *Moringa oleifera* leaf extract and its phytochemicals.




Fig 3: Schematic representation to show fertility preservation options for males

Semen cryopreservation is a well-established method for fertility preservation in adult males. However, it cannot be applicable for pre-pubertal individuals as they do not produce spermatozoa; testicular cryopreservation or SSC cryopreservation followed by derivation of spermatozoa by auto-transplantation or in vitro culture are the only available options. But these protocols are still in the experimental stages and have technical limitations. Since the survival rate in childhood cancer has increased more than 80%⁴⁷, there is a need for developing a strategy for fertility preservation, which is practically feasible and economically affordable.

Chemoprotection is considered to be an ideal, alternative option for fertility preservation using cyto-protective agents. These include Gonadotropin releasing hormone (GnRH) analogues^{48,49}, synthetic compounds like amifostine (WR-2721), AS101 (immuno-modulator)^{50,51} and natural products like antioxidants, vitamins and plant extracts.⁵²⁻⁶³ Among these, natural products have gained popularity because they are part of our diet and well tolerated by the body. Moreover, the clinical trials with GnRH analogs and other synthetic agents did not show any promising role in preventing chemotherapy-induced male gonadal toxicity.^{64,65} The chemoprotective effect of various plant extracts and natural compounds

against chemo and radiotherapy induced testicular toxicity are mentioned in Table 3.

Our earlier study demonstrated that administration of ethanolic extract of MO leaves (MOE) to pre-pubertal mice mitigated the cyclophosphamide (CP) induced testicular toxicity and improved the sperm functional characteristics when mice attained puberty.⁶⁶

The improvements in the testicular functions were associated with decreased lipid peroxidation and increased activity of antioxidant enzymes like superoxide dismutase (SOD) and catalase suggesting that MOE prevents testicular tissue by preventing oxidative stress induced by CP. Further, administration of MOE prior to CP was able to decrease the DNA damage and apoptosis in spermatogonial cells of pre-pubertal mice which are highly sensitive to CP. The protective effect of MOE is further supported by gene expression analysis by quantitative reverse transcriptase PCR (qRT-PCR), where it has been shown to modulate the expression of genes related to DNA damage response, pluripotency and stem cell survival.⁶⁷

Mechanism of protective action

High Performance Thin Layer Chromatography (HPTLC) analysis showed that MOE is rich in quercetin and chlorogenic acid that are known to possess antioxidant and anti-apoptotic properties. CP is an alkylating agent which mainly targets rapidly proliferating spermatogonial cells. In our study, it was demonstrated that MOE increased the activity of antioxidant enzymes like superoxide dismutase (SOD), catalase, glutathione peroxidase (GPX4), glutathione S transferase (GST) and glutathione reductase (GSR). These enzymes can help in CP-induced oxidative stress and prevent oxidative damage of DNA, RNA and proteins. Single cell gel electrophoresis (Comet assay) and Terminal deoxynucleotidyl transferase dUTP nick end labeling (TUNEL) assay showed that DNA damage and apoptosis induced by CP was significantly decreased by MOE administration in spermatogonial cells thereby ensuring the normal spermatogenesis and improvement in quality of spermatozoa. Further, in support of the above findings, flow cytometric analysis showed that MOE attenuated CP-induced depletion of differentiating and haploid population. Gene expression analysis showed that MOE modulated the expression of genes related to apoptosis (P53, Bax, Bcl2 and CytC), pluripotency (Oct4) and stem cell survival (c-Kit) suggesting the molecular mechanism underlying protective effect. The study on kinetics of DNA damage and repair following CP treatment in spermatogonial cells revealed that MOE administration modulated the expression of γ -H2AX (double strand break sensor) as well as RAD51 and KU80 (repair proteins). These observations further confirm that MOE can alter the CP mediated DNA damage response, thereby rescuing the spermatogonial cells leading to normal spermatogenesis (unpublished findings).

Extract/active compounds	Chemotherapeutic agent	Beneficial effect
<i>Zingiber officinale</i>	Busulfan	Increased sperm count and testosterone
<i>Ginkgo biloba</i>	Doxorubicin	Improved sperm functions, decreased oxidative stress and apoptosis
<i>Allium sativum</i>		
<i>Amaranthus viridis</i>	Cyclophosphamide	Improved sperm functions, endocrine functions, antioxidant status and decreased apoptosis
<i>Rosmarinus officinalis</i>	Etoposide	Improved spermatogenesis and antioxidant status
<i>Podophyllum hexandrum</i>	Radiation	
<i>Mentha Piperita</i>		
β -carotene	Methotrexate	Decreased oxidative damage and apoptosis
Curcumin	Cisplatin	
Vitamin C	Cisplatin	
Ellagic acid	Cisplatin Cyclophosphamide	Increased sperm quality, improved spermatogenesis and antioxidant status
Lycopene	Cyclophosphamide	

Table 3: Chemo-protective role of herbal extracts and its active principles in chemo and radiotherapy-induced testicular toxicity

Conclusion and future prospects

Administration of MOE seems to be a promising strategy to prevent chemotherapy-induced testicular toxicity. This can serve as an ideal option for pre-pubertal boys undergoing chemotherapy. The use of MO during cancer treatment is expected to have complementary effects due to the anticancer properties of the extract in addition to its normal tissue protective effect. The anti-proliferative activity of the extract helps in reducing the dose of anticancer drugs and in turn can further reduce the testicular toxicity. However, further studies are required to establish the dual role of MO as an anticancer and protective agent.

References

- 1) Leone A, Spada A, Battezzati A, Schiraldi A, Aristil J, Bertoli S. Cultivation, Genetic, Ethnopharmacology, Phytochemistry and Pharmacology of *Moringa oleifera* Leaves: An Overview. *Int J Mol Sci.* 2015;16(6):12791-835.
- 2) Anwar F, Latif S, Ashraf M, Gilani AH. *Moringa oleifera*: a food plant with multiple medicinal uses. *Phytother Res.* 2007;21(1):17-25.
- 3) Leone A, Spada A, Battezzati A, Schiraldi A, Aristil J, Bertoli S. *Moringa oleifera* Seeds and Oil: Characteristics and Uses for Human Health. *Int J Mol Sci.* 2016;17(12). pii: E2141.
- 4) Saini RK, Sivanesan I, Keum YS. Phytochemicals of *Moringa oleifera*: a review of their nutritional, therapeutic and industrial significance. *3 Biotech.* 2016;6(2):203.
- 5) Promkum C, Kupradinun P, Tuntipopipat S, Butryee C. *Asian Pac J Cancer Prev.* 2010;11(3): 627-32.
- 6) Barichella M, Pezzoli G, Faierman SA, Raspini B, Rimoldi M, Cassani E et al. Nutritional characterization of Zambian *Moringa oleifera*: acceptability and safety of short-term daily supplementation in a group of malnourished girls. *International journal of food sciences and nutrition.* 2018: 1-9.
- 7) Ramarosan RV, Valentin D, Arvisenet G. How to use local resources to fight malnutrition in Madagascar? A study combining a survey and a consumer test. *Appetite.* 2015; 95: 533-43.
- 8) Glover-Amengor M, Aryeetey R, Afari E, Nyarko A. Micronutrient composition and acceptability of *Moringa oleifera* leaf-fortified dishes by children in Ada-East district, Ghana. *Food Sci Nutr.* 2016;5(2):317-23.
- 9) Tshingani K, Donnen P, Mukumbi H, Duez P, Dramaix-Wilmet M. Impact of *Moringa oleifera* lam. Leaf powder supplementation versus nutritional counseling on the body mass index and immune response of HIV patients on antiretroviral therapy: a single-blind randomized control trial. *BMC Complement Altern Med.* 2017;17(1):420.
- 10) Jung IL. Soluble extract from *Moringa oleifera* leaves with a new anticancer activity. *PLoS One.* 2014;9(4):e95492.
- 11) Berkovich L, Earon G, Ron I, Rimmon A, Vexler A, Lev-Ari S. *Moringa Oleifera* aqueous leaf extract down-regulates nuclear factor-kappaB and increases cytotoxic effect of chemotherapy in pancreatic cancer cells. *BMC Complement Altern Med.* 2013;13:212.
- 12) Al-Asmari AK, Albalawi SM, Athar MT, Khan AQ, Al-Shahrani H, Islam M. *Moringa oleifera* as an Anti-Cancer Agent against Breast and Colorectal Cancer Cell Lines. *PLoS One.* 2015;10(8): e0135814.
- 13) Rajan TS, De Nicola GR, Iori R, Rollin P, Bramanti P, Mazzon E. Anticancer activity of glucomoringin isothiocyanate in human malignant astrocytoma cells. *Fitoterapia.* 2016;110:1-7.
- 14) Maiyo FC, Moodley R, Singh M. Cytotoxicity, Antioxidant and Apoptosis Studies of Quercetin-3-O Glucoside and 4-(beta-D-Glucopyranosyl-1->4-alpha-L-Rhamnopyranosyloxy)-Benzyl Isothiocyanate from *Moringa oleifera*. *Anticancer Agents Med Chem.* 2016;16(5):648-56.
- 15) Jung IL, Lee JH, Kang SC. A potential oral anticancer drug candidate, *Moringa oleifera* leaf extract, induces the apoptosis of human hepatocellular carcinoma cells. *Oncol Lett.* 2015;10(3):1597-1604.
- 16) Bose CK. Possible role of *Moringa oleifera* Lam. root in epithelial ovarian cancer. *MedGenMed : Medscape general medicine.* 2007;9(1):26.
- 17) Omodanisi EI, Aboua YG, Oguntibeju OO. Assessment of the Anti-Hyperglycaemic, Anti-Inflammatory and Antioxidant Activities of the Methanol Extract of *Moringa Oleifera* in Diabetes-Induced Nephrotoxic Male Wistar Rats. *Molecules.* 2017; 22(4).pii.E439.
- 18) Yassa HD, Tohamy AF. Extract of *Moringa oleifera* leaves ameliorates streptozotocin-induced Diabetes mellitus in adult rats. *Acta Histochem.* 2014;116(5):844-54.
- 19) Tang Y, Choi EJ, Han WC, Oh M, Kim J, Hwang JY et al. *Moringa oleifera* from Cambodia Ameliorates Oxidative Stress, Hyperglycemia, and Kidney Dysfunction in Type 2 Diabetic Mice. *Journal of medicinal food.* *J Med Food.* 2017;20(5):502-10.

- 20) Abd Eldaim MA, Shaban Abd Elrasoul A, Abd Elaziz SA. An aqueous extract from *Moringa oleifera* leaves ameliorates hepatotoxicity in alloxan-induced diabetic rats. *Biochem Cell Biol.* 2017;95(4):524-30.
- 21) Paula PC, Sousa DO, Oliveira JT, Carvalho AF, Alves BG, Pereira ML et al. A Protein Isolate from *Moringa oleifera* Leaves Has Hypoglycemic and Antioxidant Effects in Alloxan-Induced Diabetic Mice. *Molecules.* 2017;22.pii: E271.
- 22) Olayaki LA, Irekpita JE, Yakubu MT, Ojo OO. Methanolic extract of *Moringa oleifera* leaves improves glucose tolerance, glycogen synthesis and lipid metabolism in alloxan-induced diabetic rats. *Journal of basic and clinical physiology and pharmacology. J Basic Clin Physiol Pharmacol.* 2015;26(6):585-93.
- 23) Abd El Latif A, El Bialy Bel S, Mahboub HD, Abd Eldaim MA. *Moringa oleifera* leaf extract ameliorates alloxan-induced diabetes in rats by regeneration of beta cells and reduction of pyruvate carboxylase expression. *Biochem Cell Biol.* 2014;92(5):413-9.
- 24) Sholapur HN, Patil BM. Effect of *Moringa oleifera* bark extracts on dexamethasone-induced insulin resistance in rats. *Drug Res (Stuttg).* 2013;63(10): 527-31.
- 25) Moura MC, Trentin DS, Napoleao TH, Primon-Barros M, Xavier AS, Carneiro NP et al. Multi-effect of the water-soluble *Moringa oleifera* lectin against *Serratia marcescens* and *Bacillus* sp: antibacterial, antibiofilm and anti-adhesive properties. *Journal of applied microbiology. J Appl Microbiol.* 2017;123(4): 861-74.
- 26) Maurya SK, Singh AK. Clinical Efficacy of *Moringa oleifera* Lam. Stems Bark in Urinary Tract Infections. *Int Sch Res Notices.* 2014;2014:906843.
- 27) Feustel S, Ayon-Perez F, Sandoval-Rodriguez A, Rodríguez-Echevarría R, Contreras-Salinas H, Armendáriz-Borunda J et al. Protective Effects of *Moringa oleifera* on HBV Genotypes C and H Transiently Transfected Huh7 Cells. *J Immunol Res.* 2017;2017:6063850.
- 28) Pontual EV, Napoleao TH, Dias de Assis CR, de Souza Bezerra R, Xavier HS, Navarro DM et al. Effect of *Moringa oleifera* flower extract on larval trypsin and acetylcholinesterase activities in *Aedes aegypti*. *Arch Insect Biochem Physiol.* 2012 ;79(3): 135-52.
- 29) Chin CY, Jalil J, Ng PY, Ng SF. Development and formulation of *Moringa oleifera* standardised leaf extract film dressing for wound healing application. *J Ethnopharmacol.* 2018;212:188-99.
- 30) Rathi BS, Bodhankar SL, Baheti AM. Evaluation of aqueous leaves extract of *Moringa oleifera* Linn for wound healing in albino rats. *Indian J Exp Biol.* 2006;44(11):898-901.
- 31) Adedapo AA, Falayi OO, Oyagbemi AA. Evaluation of the analgesic, anti-inflammatory, anti-oxidant, phytochemical and toxicological properties of the methanolic leaf extract of commercially processed *Moringa oleifera* in some laboratory animals. *J Basic Clin Physiol Pharmacol.* 2015;26(5):491-9.
- 32) Gull I, Javed A, Aslam MS, Mushtaq R, Athar MA. Use of *Moringa oleifera* Flower Pod Extract as Natural Preservative and Development of SCAR Marker for Its DNA Based Identification. *Biomed Res Int.* 2016;2016:7584318.
- 33) Jaja-Chimedza A, Graf BL, Simmler C, Kim Y, Kuhn P, Pauli GF et al. Biochemical characterization and anti-inflammatory properties of an isothiocyanate-enriched moringa (*Moringa oleifera*) seed extract. *PLoS One.* 2017;12(8): e0182658 658.
- 34) Faizi S, Siddiqui BS, Saleem R, Aftab K, Shaheen F, Gilani AH. Hypotensive constituents from the pods of *Moringa oleifera*. *Planta Med.* 1998; 64(3):225-8.
- 35) Kim Y, Wu AG, Jaja-Chimedza A, Graf BL, Waterman C, Verzi MP et al. Isothiocyanate-enriched moringa seed extract alleviates ulcerative colitis symptoms in mice. *PLoS One.* 2017;12(9): e0184709.
- 36) Kirisattayakul W, Wattanathorn J, Tong-Un T, Muchimapura S, Wannanon P, Jittiwat J. Cerebroprotective effect of *Moringa oleifera* against focal ischemic stroke induced by middle cerebral artery occlusion. *Oxid Med Cell Longev.* 2013;2013: 951415.
- 37) Ray K, Hazra R, Guha D. Central inhibitory effect of *Moringa oleifera* root extract: possible role of neurotransmitters. *Indian J Exp Biol.* 2003;41(11): 1279-84.
- 38) Sinha M, Das DK, Bhattacharjee S, Majumdar S, Dey S. Leaf extract of *Moringa oleifera* prevents ionizing radiation-induced oxidative stress in mice. *J Med Food.* 2011;14(10):1167-72.
- 39) Rao AV, Devi PU, Kamath R. In vivo radioprotective effect of *Moringa oleifera* leaves. *Indian J Exp Biol.* 2001 Sep;39(9):858-63.
- 40) Igado OO, Glaser J, Ramos-Tirado M, Bankoğlu EE, Atiba FA, Holzgrabe U et al. Isolation of a novel compound (MIMO₂) from the methanolic extract of *Moringa oleifera* leaves: protective effects against vanadium-induced cytotoxicity. *Drug Chem Toxicol.* 2018;41(3):249-58.

- 41) Karthivashan G, Kura AU, Arulsevan P, Md Isa N, Fakurazi S. The modulatory effect of *Moringa oleifera* leaf extract on endogenous antioxidant systems and inflammatory markers in an acetaminophen-induced nephrotoxic mice model. *PeerJ*. 2016;4:e2127.
- 42) Fakurazi S, Hairuszah I, Nanthini U. *Moringa oleifera* Lam prevents acetaminophen induced liver injury through restoration of glutathione level. *Food Chem Toxicol*. 2008;46(8):2611-5.
- 43) Pari L, Kumar NA. Hepatoprotective activity of *Moringa oleifera* on antitubercular drug-induced liver damage in rats. *Journal of medicinal food. J Med Food*. 2002;5(3):171-7.
- 44) Al-Said MS, Mothana RA, Al-Yahya MA, Al-Blowi AS, Al-Sohaibani M, Ahmed AF et al. Edible oils for liver protection: hepatoprotective potentiality of *Moringa oleifera* seed oil against chemical-induced hepatitis in rats. *J Food Sci*. 2012;77(7):T124-30.
- 45) Sadek KM. Chemotherapeutic efficacy of an ethanolic *Moringa oleifera* leaf extract against chromium-induced testicular toxicity in rats. *Andrologia*. 2014;46(9):1047-54.
- 46) Abarikwu SO, Benjamin S, Ebah SG, Obilor G, Agbam G. Oral administration of *Moringa oleifera* oil but not coconut oil prevents mercury-induced testicular toxicity in rats. *Andrologia*. 2017; 49(1).
- 47) Ward E, DeSantis C, Robbins A, Kohler B, Jemal A. Childhood and adolescent cancer statistics, 2014. *CA Cancer J Clin*. 2014;64(2):83-103.
- 48) Meistrich ML, Wilson G, Huhtaniemi I. Hormonal treatment after cytotoxic therapy stimulates recovery of spermatogenesis. *Cancer Res*. 1999; 59(15):3557-60.
- 49) Wang G, Shao SH, Weng CC, Wei C, Meistrich ML. Hormonal suppression restores fertility in irradiated mice from both endogenous and donor-derived stem spermatogonia. *Toxicol Sci*. 2010;117(1):225-37.
- 50) Vendramini V, Sasso-Cerri E, Miraglia SM. Amifostine reduces the seminiferous epithelium damage in doxorubicin-treated prepubertal rats without improving the fertility status. *Reprod Biol Endocrinol*. 2010;8:3.
- 51) Carmely A, Meirow D, Peretz A, Albeck M, Bartoov B, Sredni B. Protective effect of the immunomodulator AS101 against cyclophosphamide-induced testicular damage in mice. *Hum Reprod*. 2009;24(6):1322-9.
- 52) Vardi N, Parlakpinar H, Ates B, Cetin A, Otlu A. Antiapoptotic and antioxidant effects of beta-carotene against methotrexate-induced testicular injury. *Fertil Steril*. 2009;92(6):2028-33.
- 53) Ahmed EA, Omar HM, elghaffar S, Ragb SM, Nasser AY. The antioxidant activity of vitamin C, DPPD and L-cysteine against Cisplatin-induced testicular oxidative damage in rats. *Food Chem Toxicol*. 2011;49(5):1115-21.
- 54) Bordbar H, Esmailpour T, Dehghani F, Panjehshahin MR. Stereological study of the effect of ginger's alcoholic extract on the testis in busulfan-induced infertility in rats. *Iran J Reprod Med*. 2013;11(6):467-72.
- 55) Yeh YC, Liu TJ, Wang LC, Lee HW, Ting CT, Lee WL et al. A standardized extract of Ginkgo biloba suppresses doxorubicin-induced oxidative stress and p53-mediated mitochondrial apoptosis in rat testes. *Br J Pharmacol*. 2009;156(1):48-61.
- 56) Nasr AY. The impact of aged garlic extract on adriamycin-induced testicular changes in adult male Wistar rats. *Acta Histochem*. 2017 Jul;119(6):648-662.
- 57) Ayoka OA, Ojo OE, Imafidon EC, Ademoye KA, Oladele AA. Neuro-endocrine effects of aqueous extract of *Amaranthus viridis* (Linn.) leaf in male Wistar rat model of cyclophosphamide-induced reproductive toxicity. *Toxicol Rep*. 2016;3:608-19.
- 58) Tousson E, Bayomy MF, Ahmed AA. Rosemary extract modulates fertility potential, DNA fragmentation, injury, KI67 and P53 alterations induced by etoposide in rat testes. *Biomed Pharmacother*. 2018;98:769-74.
- 59) Samanta N, Goel HC. Protection against radiation induced damage to spermatogenesis by *Podophyllum hexandrum*. *J Ethnopharmacol*. 2002;81(2): 217-24.
- 60) Samarth RM, Samarth M. Protection against radiation-induced testicular damage in Swiss albino mice by *Mentha piperita* (Linn.). *Basic Clin Pharmacol Toxicol*. 2009;104(4):329-34.
- 61) Ilbey YO, Ozbek E, Cekmen M, Simsek A, Otunc-tumur A, Somay A. Protective effect of curcumin in cisplatin-induced oxidative injury in rat testis: mitogen-activated protein kinase and nuclear factor-kappa B signaling pathways. *Hum Reprod*. 2009;24(7):1717-25.

- 62) Turk G, Atessahin A, Sonmez M, Ceribasi AO, Yuce A. Improvement of cisplatin-induced injuries to sperm quality, the oxidant-antioxidant system, and the histologic structure of the rat testis by ellagic acid. *Fertil Steril.* 2008;89(5 Suppl): 1474-81.
- 63) Ceribasi AO, Turk G, Sonmez M, Sakin F, Atessahin A. Toxic effect of cyclophosphamide on sperm morphology, testicular histology and blood oxidant-antioxidant balance, and protective roles of lycopene and ellagic acid. *Basic Clin Pharmacol Toxicol.* 2010;107(3):730-6.
- 64) da Cunha MF, Meistrich ML, Nader S. Absence of testicular protection by a gonadotropin-releasing hormone analogue against cyclophosphamide-induced testicular cytotoxicity in the mouse. *Cancer Res.* 1987;47(4):1093-7.
- 65) Vendramini V, Robaire B, Miraglia SM. Amifostine-doxorubicin association causes long-term prepubertal spermatogonia DNA damage and early developmental arrest. *Hum Reprod.* 2012;27(8):2457-66.
- 66) Nayak G, Vadinkar A, Nair S, Kalthur SG, D'Souza AS, Shetty PK et al. Sperm abnormalities induced by pre-pubertal exposure to cyclophosphamide are effectively mitigated by *Moringa oleifera* leaf extract. *Andrologia.* 2016;48(2):125-36.
- 67) Nayak G, Honguntikar SD, Kalthur SG, D'Souza AS, Mutalik S, Setty MM et al. Ethanolic extract of *Moringa oleifera* Lam. leaves protect the pre-pubertal spermatogonial cells from cyclophosphamide-induced damage. *J Ethnopharmacol.* 2016;182:101-9.